

Chapter 5A: Continuous Random Variables, PDFs, and Uniform Distribution

MATH/STAT 394: Probability I

Arman Jahangiri

University of Washington
Department of Mathematics

Summer 2026

By the end of this lecture, students should be able to:

- ▶ define and interpret PDFs and CDFs;
- ▶ compute probabilities by integrating a PDF;
- ▶ recognize valid PDFs;
- ▶ compute expectation for continuous random variables;
- ▶ define and use the Uniform distribution;
- ▶ understand the universality of the Uniform distribution.

Continuous Random Variables

From Discrete to Continuous Random Variables

Discrete random variables take values in a finite or countably infinite set.

Continuous random variables can take any value in an interval, such as

$$(0, 1), \quad (a, b), \quad [0, \infty), \quad \mathbb{R}.$$

Main difference

For a continuous random variable,

$$\mathbb{P}(X = x) = 0$$

for every individual point x .

Definition

For a continuous random variable X , the probability density function, or PDF, is a function f such that

$$\mathbb{P}(X \in A) = \int_A f(x) dx, \quad \forall A \subseteq \mathbb{R}$$

The value $f(x)$ itself is not a probability (could be greater than 1).

Probabilities are areas under the density curve.

How to verify a function is a PDF?

A function f is a valid PDF if it satisfies two properties, namely

$$f(x) \geq 0 \quad \text{for all } x,$$

and

$$\int_{-\infty}^{\infty} f(x) dx = 1.$$

Analogy

For discrete random variables:

$$p_X(x) \geq 0, \quad \sum_x p_X(x) = 1.$$

For continuous random variables:

$$f_X(x) \geq 0, \quad \int_{-\infty}^{\infty} f(x) dx = 1.$$

CDF has the same formula for continuous and discrete r.v.s, i.e.

$$F(x) = \mathbb{P}(X \leq x).$$

For a continuous random variable with PDF f , according to the definition of PDF,

$$F(x) = \int_{-\infty}^x f(t) dt.$$

Note that , according to the definition of CDF,

$$\mathbb{P}(a < X \leq b) = F(b) - F(a)$$

This, together with the Fundamental Theorem of Calculus, implies that

$$\frac{d}{dx} F(x) = F'(x) = f(x) \tag{1}$$

For **continuous** r.v.s, since $\mathbb{P}(X = a) = 0$ for any $a \in \mathbb{R}$,

$$\mathbb{P}(X \leq a) = \mathbb{P}(X < a),$$

$$\mathbb{P}(X \geq a) = \mathbb{P}(X > a),$$

$$\mathbb{P}(a \leq X \leq b) = \mathbb{P}(a < X \leq b) = \mathbb{P}(a \leq X < b) = \mathbb{P}(a < X < b).$$

What Does the Density $f(x)$ Mean?

For a continuous random variable,

$$F(x) = \mathbb{P}(X \leq x) = \int_{-\infty}^x f(t) dt.$$

For a small number $\epsilon > 0$,

$$\mathbb{P}(x \leq X \leq x + \epsilon) = F(x + \epsilon) - F(x).$$

Therefore,

$$\frac{\mathbb{P}(x \leq X \leq x + \epsilon)}{\epsilon} = \frac{F(x + \epsilon) - F(x)}{\epsilon}.$$

Taking the limit as $\epsilon \rightarrow 0$,

$$\lim_{\epsilon \rightarrow 0} \frac{\mathbb{P}(x \leq X \leq x + \epsilon)}{\epsilon} = \lim_{\epsilon \rightarrow 0} \frac{F(x + \epsilon) - F(x)}{\epsilon}.$$

By the definition of derivative,

$$f(x) = F'(x) = \lim_{\epsilon \rightarrow 0} \frac{F(x + \epsilon) - F(x)}{\epsilon}.$$

Thus,

$$f(x) = \lim_{\epsilon \rightarrow 0} \frac{\mathbb{P}(x \leq X \leq x + \epsilon)}{\epsilon}$$

Definition

If X is continuous with PDF f , then

$$\mathbb{E}(X) = \int_{-\infty}^{\infty} xf(x) dx.$$

If the support is an interval, we integrate over that interval.

Expectation is still the balancing point of the distribution.

LOTUS

If X has PDF f and $g : \mathbb{R} \rightarrow \mathbb{R}$, then

$$\mathbb{E}(g(X)) = \int_{-\infty}^{\infty} g(x)f(x) dx.$$

Proof idea. For a small interval $[x, x + \Delta x]$,

$$\mathbb{P}(x \leq X \leq x + \Delta x) \approx f(x)\Delta x.$$

On this interval, $g(X)$ is approximately $g(x)$. Hence,

$$g(X)\mathbb{P}(x \leq X \leq x + \Delta x) \approx g(x)f(x)\Delta x.$$

Adding over many small intervals,

$$\mathbb{E}(g(X)) \approx \sum_i g(x_i)f(x_i)\Delta x.$$

Taking the limit as $\Delta x \rightarrow 0$, the Riemann sum becomes

$$\mathbb{E}(g(X)) = \int_{-\infty}^{\infty} g(x)f(x) dx.$$

Uniform Distribution

Definition

A random variable U has the Uniform distribution on (a, b) if its PDF is

$$f(x) = \begin{cases} \frac{1}{b-a}, & a < x < b, \\ 0, & \text{otherwise.} \end{cases}$$

We write

$$U \sim \text{Unif}(a, b).$$

PDF Verification

$$f(x) = \frac{1}{b-a} > 0$$

since $b > a$. And

$$\int_{-\infty}^{\infty} f(x) dx = \int_a^b \frac{1}{b-a} dx = \frac{1}{b-a}(b-a) = 1.$$

If

$$U \sim \text{Unif}(a, b),$$

then its CDF is

$$F(x) = \begin{cases} 0, & x \leq a, \\ \frac{x-a}{b-a}, & a < x < b, \\ 1, & x \geq b. \end{cases}$$

For the standard Uniform,

$$U \sim \text{Unif}(0, 1),$$

we have

$$F(x) = x, \quad 0 < x < 1.$$

Universality of the Uniform

Why the Uniform Distribution Is Special

The Uniform distribution is a universal starting point for simulation.

Many algorithms first generate

$$U \sim \text{Unif}(0, 1),$$

then transform U to obtain another distribution.

Main idea

A Uniform random variable can be transformed into a random variable with many other distributions.

Suppose F is a continuous, strictly increasing CDF.

Let

$$U \sim \text{Unif}(0, 1).$$

Define

$$X = F^{-1}(U).$$

Then

X has CDF F .

This is called the inverse CDF method or inverse transform sampling.

Let

$$X = F^{-1}(U).$$

Then

$$\mathbb{P}(X \leq x) = \mathbb{P}(F^{-1}(U) \leq x).$$

Since F is increasing,

$$F^{-1}(U) \leq x \iff U \leq F(x).$$

Therefore,

$$\mathbb{P}(X \leq x) = \mathbb{P}(U \leq F(x)).$$

Because $U \sim \text{Unif}(0, 1)$,

$$\mathbb{P}(U \leq F(x)) = F(x).$$

Thus X has CDF F .

Conversely, if X has continuous strictly increasing CDF F , then

$$F(X) \sim \text{Unif}(0, 1).$$

Interpretation

Applying the CDF to a random variable converts its value into a percentile.

For example, if exam scores have CDF F , then $F(X)$ is the percentile of a randomly selected student.

Uniform Mean and Variance

If $U \sim \text{Unif}(a, b)$, then

$$f_U(u) = \frac{1}{b-a}, \quad a \leq u \leq b.$$

Mean

$$\mathbb{E}(U) = \int_a^b u \frac{1}{b-a} du = \frac{1}{b-a} \left[\frac{u^2}{2} \right]_a^b = \frac{b^2 - a^2}{2(b-a)} = \frac{a+b}{2}.$$

Variance

First,

$$\mathbb{E}(U^2) = \int_a^b u^2 \frac{1}{b-a} du = \frac{b^3 - a^3}{3(b-a)} = \frac{a^2 + ab + b^2}{3}.$$

Thus,

$$\text{Var}(U) = \mathbb{E}(U^2) - [\mathbb{E}(U)]^2 = \frac{a^2 + ab + b^2}{3} - \left(\frac{a+b}{2} \right)^2 = \frac{(b-a)^2}{12}.$$

If

$$U \sim \text{Unif}(0, 1),$$

then

$$a + (b - a)U \sim \text{Unif}(a, b).$$

This transforms:

$$(0, 1) \longrightarrow (a, b).$$

Useful idea

Start with the standard Uniform and shift/scale to get any Uniform interval.

Summary

- ▶ Continuous random variables have probabilities over intervals, not points.
- ▶ A PDF integrates to probabilities:

$$\mathbb{P}(a < X < b) = \int_a^b f(x) dx.$$

- ▶ A valid PDF is nonnegative and integrates to 1.
- ▶ The CDF is accumulated area:

$$F(x) = \int_{-\infty}^x f(t) dt.$$

- ▶ Uniform probabilities are proportional to interval length.
- ▶ The Uniform distribution is universal through the inverse CDF method.

Next lecture

Normal and Exponential distributions.

Temporary page!

$\text{\LaTeX}$ was unable to guess the total number of pages correctly. As there was some unprocessed data have been added to the final page this extra page has been added to receive it.
If you rerun the document (without altering it) this surplus page will go away, because $\text{\LaTeX}$ now knows the number of pages to expect for this document.