

Chapter 6B: Moment Generating Functions

MATH/STAT 394: Probability I

Arman Jahangiri

University of Washington Department of Mathematics

Summer 2026

By the end of this lecture, students should be able to:

- ▶ define the moment generating function (MGF);
- ▶ compute MGFs of common distributions;
- ▶ extract moments from MGFs;
- ▶ understand how MGFs determine distributions;
- ▶ compute distributions of sums using MGFs;
- ▶ derive moments of Exponential and Normal distributions using MGFs.

Definition

The moment generating function (MGF) of X is

$$M_X(t) = \mathbb{E}(e^{tX})$$

provided the expectation is finite in some open interval around 0.

Equivalently,

$$M_X(t) = \begin{cases} \sum_x e^{tx} p_X(x), & \text{if } X \text{ is discrete,} \\ \int_{-\infty}^{\infty} e^{tx} f_X(x) dx, & \text{if } X \text{ is continuous.} \end{cases}$$

Quick check

Every valid MGF satisfies

$$M_X(0) = 1.$$

$$M_X(t) = \mathbb{E}(e^{tX}) = \sum_x e^{tx} p_X(x).$$

Bernoulli

If $X \sim \text{Bern}(p)$, then

$$M_X(t) = e^{t \cdot 1} p + e^{t \cdot 0} (1 - p) = pe^t + q.$$

Discrete Uniform

If $X \sim \text{DUnif}\{1, \dots, m\}$, then

$$M_X(t) = \frac{1}{m} \sum_{k=1}^m e^{tk} = \frac{1}{m} (e^t + e^{2t} + \dots + e^{mt}) = \frac{e^t(e^{mt} - 1)}{m(e^t - 1)}.$$

Binomial

If $X \sim \text{Bin}(n, p)$, then

$$M_X(t) = \sum_{k=0}^n e^{tk} \binom{n}{k} p^k q^{n-k} = \sum_{k=0}^n \binom{n}{k} (pe^t)^k q^{n-k} = (q + pe^t)^n.$$

Poisson

If $X \sim \text{Pois}(\lambda)$, then

$$M_X(t) = \sum_{k=0}^{\infty} e^{tk} e^{-\lambda} \frac{\lambda^k}{k!} = e^{-\lambda} \sum_{k=0}^{\infty} \frac{(\lambda e^t)^k}{k!} = e^{-\lambda} e^{\lambda e^t} = e^{\lambda(e^t - 1)}.$$

Geometric

If $X \sim \text{Geom}(p)$, where X counts failures before the first success, then

$$M_X(t) = \sum_{k=0}^{\infty} e^{tk} q^k p = p \sum_{k=0}^{\infty} (qe^t)^k = \frac{p}{1 - qe^t}, \quad qe^t < 1.$$

Negative Binomial

If $X \sim \text{NegBin}(r, p)$, where X counts failures before the r -th success, then

$$M_X(t) = \sum_{k=0}^{\infty} e^{tk} \binom{k+r-1}{k} p^r q^k = p^r \sum_{k=0}^{\infty} \binom{k+r-1}{k} (qe^t)^k = \left(\frac{p}{1 - qe^t} \right)^r.$$

Hypergeometric

If $X \sim \text{HGeom}(w, b, n)$,

then

$$\mathbb{P}(X = k) = \frac{\binom{w}{k} \binom{b}{n-k}}{\binom{w+b}{n}},$$

where

$$\max(0, n - b) \leq k \leq \min(w, n).$$

Therefore,

$$M_X(t) = \sum_{k=\max(0, n-b)}^{\min(w, n)} e^{tk} \frac{\binom{w}{k} \binom{b}{n-k}}{\binom{w+b}{n}}.$$

Note: Unlike Binomial or Poisson, this MGF usually does not simplify to a nice elementary closed form.

$$M_X(t) = \mathbb{E}(e^{tX}) = \int_{-\infty}^{\infty} e^{tx} f_X(x) dx.$$

Uniform

If $X \sim \text{Unif}(a, b)$, then

$$M_X(t) = \int_a^b e^{tx} \frac{1}{b-a} dx = \frac{1}{b-a} \left[\frac{e^{tx}}{t} \right]_a^b = \frac{e^{tb} - e^{ta}}{t(b-a)}, \quad t \neq 0.$$

Also, $M_X(0) = 1$.

Exponential

If $X \sim \text{Exp}(\lambda)$, then

$$M_X(t) = \int_0^{\infty} e^{tx} \lambda e^{-\lambda x} dx = \lambda \int_0^{\infty} e^{-(\lambda-t)x} dx = \frac{\lambda}{\lambda-t}, \quad t < \lambda.$$

Gamma

If $X \sim \text{Gamma}(\alpha, \lambda)$, with shape α and rate λ , then

$$M_X(t) = \int_0^{\infty} e^{tx} \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} dx = \frac{\lambda^\alpha}{\Gamma(\alpha)} \int_0^{\infty} x^{\alpha-1} e^{-(\lambda-t)x} dx.$$

Let $u = (\lambda - t)x$. Then $x = u/(\lambda - t)$, $dx = du/(\lambda - t)$, and

$$M_X(t) = \frac{\lambda^\alpha}{\Gamma(\alpha)} \frac{1}{(\lambda - t)^\alpha} \underbrace{\int_0^{\infty} u^{\alpha-1} e^{-u} du}_{\Gamma(\alpha)} = \frac{\lambda^\alpha}{(\lambda - t)^\alpha} = \left(\frac{\lambda}{\lambda - t} \right)^\alpha, \quad t < \lambda.$$

Standard Normal

Let $Z \sim N(0, 1)$. Then

$$M_Z(t) = \mathbb{E}(e^{tZ}) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{tz} e^{-z^2/2} dz.$$

Complete the square:

$$tz - \frac{z^2}{2} = -\frac{1}{2}(z^2 - 2tz) = -\frac{1}{2}((z - t)^2 - t^2) = -\frac{(z - t)^2}{2} + \frac{t^2}{2}.$$

Therefore,

$$M_Z(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-(z-t)^2/2} e^{t^2/2} dz = e^{t^2/2} \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-(z-t)^2/2} dz.$$

Since the last integral equals $\sqrt{2\pi}$,

$$M_Z(t) = e^{t^2/2}.$$

Normal

Let $X \sim N(\mu, \sigma^2)$. Write

$$X = \mu + \sigma Z, \quad Z \sim N(0, 1).$$

Then

$$M_X(t) = \mathbb{E}(e^{tX}) = \mathbb{E}(e^{t(\mu + \sigma Z)}) = e^{\mu t} \mathbb{E}(e^{\sigma t Z}).$$

Using

$$M_Z(s) = e^{s^2/2},$$

with $s = \sigma t$, we get

$$M_X(t) = e^{\mu t} M_Z(\sigma t) = e^{\mu t} e^{\sigma^2 t^2 / 2} = e^{\mu t + \frac{1}{2} \sigma^2 t^2}.$$

Moments from MGFs

Theorem

The n^{th} population moment is obtained by differentiating the MGF:

$$\mathbb{E}(X^n) = M_X^{(n)}(0).$$

Examples:

$$\mathbb{E}(X) = M_X'(0), \quad \mathbb{E}(X^2) = M_X''(0).$$

Proof: MGFs Generate Moments

Recall that the moment generating function is

$$M_X(t) = \mathbb{E}(e^{tX}).$$

Differentiate once:

$$M'_X(t) = \frac{d}{dt} \mathbb{E}(e^{tX}) = \mathbb{E} \left(\frac{d}{dt} e^{tX} \right) = \mathbb{E}(Xe^{tX}).$$

Now plugging in $t = 0$,

$$M'_X(0) = \mathbb{E}(Xe^0) = \mathbb{E}(X).$$

Differentiate another time noting that X is a constant w.r.t. t .

$$M''_X(t) = \frac{d}{dt} \mathbb{E}(Xe^{tX}) = \mathbb{E} \left(\frac{d}{dt} Xe^{tX} \right) = \mathbb{E}(X^2 e^{tX}).$$

Plugging in $t = 0$,

$$M''_X(0) = \mathbb{E}(X^2 e^0) = \mathbb{E}(X^2).$$

Similarly, differentiating n times gives

$$M_X^{(n)}(t) = \mathbb{E}(X^n e^{tX}).$$

Therefore, at $t = 0$,

$$M_X^{(n)}(0) = \mathbb{E}(X^n e^0) = \mathbb{E}(X^n).$$

The Taylor expansion of

$$e^{tX}$$

is

$$e^{tX} = 1 + tX + \frac{t^2 X^2}{2!} + \dots .$$

Taking expectations:

$$M_X(t) = 1 + t\mathbb{E}(X) + \frac{t^2}{2!}\mathbb{E}(X^2) + \dots .$$

The moments appear as Taylor coefficients.

Example: Bernoulli Moments

For

$$M_X(t) = pe^t + q,$$

we compute:

$$M'_X(t) = pe^t.$$

Thus:

$$M'_X(0) = p.$$

So

$$\mathbb{E}(X) = p.$$

Theorem (Uniqueness of MGF)

If two random variables have the same MGF on an interval around 0, then they have the same distribution, i.e., the moment generating function uniquely determines the distribution.

Thus MGFs characterize distributions, similarly to PDFs and CDFs.

MGFs of Sums

Theorem

If X and Y are independent, then

$$M_{X+Y}(t) = M_X(t) \times M_Y(t).$$

More generally, if $X_1, \dots, X_n$ are independent, then

$$M_{\sum_{i=1}^n X_i}(t) = \prod_{i=1}^n M_{X_i}(t)$$

This is one of the main reasons MGFs are useful. Since using the MGF Uniqueness theorem, the MGF characterizes the distribution in full.

A Binomial random variable can be written as:

$$X = I_1 + \cdots + I_n,$$

where the I_j are independent Bernoulli(p) random variables.

Since each Bernoulli MGF is

$$pe^t + q,$$

we get

$$M_X(t) = (pe^t + q)^n.$$

If

$$X \sim \text{NBin}(r, p),$$

then

$$X = G_1 + \cdots + G_r,$$

where the G_j are independent $\text{Geometric}(p)$ random variables.

Thus,

$$M_X(t) = \left(\frac{p}{1 - qe^t} \right)^r.$$

Sums via MGFs

If

$$X \sim \text{Pois}(\lambda), \quad Y \sim \text{Pois}(\mu),$$

independently, then

$$M_X(t) = e^{\lambda(e^t - 1)},$$

and

$$M_Y(t) = e^{\mu(e^t - 1)}.$$

Therefore,

$$M_{X+Y}(t) = e^{(\lambda+\mu)(e^t - 1)}.$$

Hence:

$$X + Y \sim \text{Pois}(\lambda + \mu).$$

Remark: This can be generalized to n independent Poisson r.v.s using induction.

If

$$X_1 \sim N(\mu_1, \sigma_1^2), \quad X_2 \sim N(\mu_2, \sigma_2^2),$$

independently, then

$$M_{X_1+X_2}(t) = e^{(\mu_1+\mu_2)t + \frac{1}{2}(\sigma_1^2+\sigma_2^2)t^2}.$$

Thus,

$$X_1 + X_2 \sim N(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2).$$

Remark: This can be generalized to n independent Normal r.v.s using induction.

Distributions Closed Under Addition Under Independence

Distribution	Assumption	Sum Distribution
Bernoulli	$X_i \sim \text{Bern}(p)$	$\sum_{i=1}^n X_i \sim \text{Bin}(n, p)$
Binomial	$X_i \sim \text{Bin}(n_i, p)$	$\sum_{i=1}^n X_i \sim \text{Bin}\left(\sum n_i, p\right)$
Poisson	$X_i \sim \text{Pois}(\lambda_i)$	$\sum_{i=1}^n X_i \sim \text{Pois}\left(\sum \lambda_i\right)$
Negative Binomial	$X_i \sim \text{NegBin}(r_i, p)$	$\sum_{i=1}^n X_i \sim \text{NegBin}\left(\sum r_i, p\right)$
Geometric	$X_i \sim \text{Geom}(p)$	$\sum_{i=1}^n X_i \sim \text{NegBin}(n, p)$
Normal	$X_i \sim N(\mu_i, \sigma_i^2)$	$\sum_{i=1}^n X_i \sim N\left(\sum \mu_i, \sum \sigma_i^2\right)$
Gamma	$X_i \sim \text{Gamma}(\alpha_i, \lambda)$	$\sum_{i=1}^n X_i \sim \text{Gamma}\left(\sum \alpha_i, \lambda\right)$
Exponential	$X_i \sim \text{Exp}(\lambda)$	$\sum_{i=1}^n X_i \sim \text{Gamma}(n, \lambda)$

Summary

- ▶ MGFs encode moments through derivatives.
- ▶ MGFs determine distributions.
- ▶ Independent sums correspond to products of MGFs.
- ▶ MGFs simplify moment calculations dramatically.
- ▶ The Normal and Exponential MGFs are especially important.

Next lecture

Joint distributions.